

chain nodes :

7 8

ring nodes :

1 2 3 4 5 6 9 10 11 12 13 14 15 16 17 18 19 20 21 22

chain bonds :

5-7 7-8 8-9

ring bonds :

1-2 1-6 2-3 2-19 3-4 3-22 4-5 5-6 9-10 9-14 10-11 11-12 11-15  
12-13 12-18 13-14 15-16 16-17 17-18 19-20 20-21 21-22

exact bonds :

5-7 7-8 8-9

normalized bonds :

1-2 1-6 2-3 2-19 3-4 3-22 4-5 5-6 9-10 9-14 10-11 11-12 11-15  
12-13 12-18 13-14 15-16 16-17 17-18 19-20 20-21 21-22

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS8:CLASS9:Atom  
10:Atom 11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom  
19:Atom 20:Atom 21:Atom 22:Atom

SAMPLE SEARCH INITIATED 12:22:39 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 124 TO ITERATE

100.0% PROCESSED 124 ITERATIONS 2  
ANSWERS  
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*  
PROJECTED ITERATIONS: 1812 TO 3148  
PROJECTED ANSWERS: 2 TO 124

L3 2 SEA SSS SAM L1

L4 2 L3

=> D L4 IBIB ABS HITSTR 1-2

L4 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2004:789773 CAPLUS  
DOCUMENT NUMBER: 142:48473  
TITLE: Inhibition of human immunodeficiency virus  
type I  
integrase by naphthamidines and  
2-aminobenzimidazoles  
AUTHOR(S): Middleton, Tim; Lim, Hock B.; Montgomery,  
Debra;  
Rockway, Todd; Tang, Hua; Cheng, Xueheng; Lu,  
Liangjun; Mo, Hongmei; Kohlbrenner, William  
E.; Molla,  
Akhteruzzaman; Kati, Warren M.  
CORPORATE SOURCE: Global Pharmaceutical Research and  
Development,  
Department R47D, Abbott Laboratories, Abbott  
Park, IL,  
60064-6217, USA  
SOURCE: Antiviral Research (2004), 64(1), 35-45  
CODEN: ARSRDR; ISSN: 0166-3542  
PUBLISHER: Elsevier B.V.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Retroviral integrases catalyze two of the steps of insertion of  
proviral  
DNA into the host genomic DNA. Inhibitors that target the  
second step,  
strand transfer into the host DNA, have been demonstrated to have  
antiviral activity in cell culture. The authors describe two  
classes of

HIV-1 integrase inhibitors that block strand transfer, one based on a naphthamidine core and one on a benzimidazole core. While the naphthamidine compds. showed some propensity to interact with the DNA

substrate, both classes were shown to bind directly to integrase. The

naphthamidine compds. showed activity in cell culture, and a direct effect

on integrase was indicated by an increase in 2-LTR products in the

presence of a naphthamidine compound These two classes of compds. represent

potential starting points for the development of new classes of integrase inhibitors.

IT 808144-92-5

RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic

use); BIOL (Biological study); USES (Uses)


(inhibition of human immunodeficiency virus type I integrase by

naphthamidines and 2-aminobenzimidazoles)

RN 808144-92-5 CAPLUS

CN 2-Naphthalenecarboximidamide,

6-[(7-phenyl-2-naphthalenyl)ethynyl]- (9CI)  
(CA INDEX NAME)



REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES  
AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE

RE FORMAT

L4 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:55291 CAPLUS


DOCUMENT NUMBER: 140:199098

TITLE: Synthesis and properties of monosubstituted ethynylcorannulenes

AUTHOR(S): Jones, Carissa S.; Elliott, Eric; Siegel, Jay S.

CORPORATE SOURCE: Department of Chemistry, University of California, San

SOURCE: Diego, La Jolla, CA, 92093-0358, USA  
 Synlett (2004), (1), 187-191  
 CODEN: SYNLES; ISSN: 0936-5214  
 PUBLISHER: Georg Thieme Verlag  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 140:199098  
 GI




I


AB The solution-phase synthesis of corannulene (I) has been modified and it is now possible to prepare multi gram quantities of corannulene more efficiently, with considerably less toxic reagents.

Cross-coupling of bromocorannulene with TMS-acetylene and phenylacetylene affords novel ethynyl-containing corannulene derivs. Deprotection of TMS-ethynyl corannulene affords the naked alkyne, which can be cross-coupled with pentafluoriodobenzene and bromocorannulene to afford the appropriate alkyne derivs. The photophys. properties of this new and novel family of alkyne-containing corannulene derivs. has been evaluated and all of the new derivs. exhibit low to moderate quantum efficiencies.

IT 663617-09-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of arylethynylcorannulene and bicorannulenylacetylene via cross coupling of ethynylcorannulene with pentafluoriodobenzene and bromocorannulene)

RN 663617-09-2 CAPLUS  
 CN Dibenzo[ghi,mno]fluoranthene, 1,1'-(1,2-ethynediyl)bis- (9CI)  
 (CA INDEX NAME)





chain nodes :

1 2 3 4

chain bonds :

1-2 1-4 2-3

exact/norm bonds :

1-4 2-3

exact bonds :

1-2

Match level :

1:CLASS2:CLASS3:Atom 4:Atom

L4 50 SEA SSS SAM L1

L5 45 L4

=> S L5 AND SEMICONDUCTOR

502507 SEMICONDUCTOR

L6 2 L5 AND SEMICONDUCTOR

=> D L6 IBIB ABS HITSTR 1-2

L6 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:512808 CAPLUS

DOCUMENT NUMBER: 141:80542

TITLE: Organic semiconductive materials and  
manufacture of

organic electric device by film formation of  
the materials

INVENTOR(S): Takada, Yoshihiro; Aramaki, Shinji

PATENT ASSIGNEE(S): Mitsubishi Chemical Corp., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 23 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.
DATE			
-----	----	-----	-----
-----			

JP 2004179249	A2	20040624	JP 2002-341175
---------------	----	----------	----------------

20021125

PRIORITY APPLN. INFO.: JP 2002-341175

20021125

AB The organic semiconductive materials comprise compds. with mol.  
weight

≤2000 and containing ≥1 structure where (substituted) aromatic  
hydrocarbon ring-containing groups or (substituted) aromatic  
heterocyclic

ring-containing groups are bonded via alkynylene group. The  
organic elec. device

such as a field-effect transistor, IC, a display, etc., is  
prepared by

film-formation of an organic semiconductives containing the  
above-mentioned

materials, heating the prepared film until it becomes a  
fluidizing phase,

preferably a liquid crystalline phase, and cooling.

IT 710338-94-6P

RL: DEV (Device component use); IMF (Industrial manufacture);


PREP

(Preparation); USES (Uses)

(organic semiconductive materials for manufacture of  
field-effect transistor by  
film formation)

RN 710338-94-6 CAPLUS

CN 2,2':5',2''-Terthiophene, 5,5''-bis[(4-butyl-2,3,5,6-  
tetrafluorophenyl)ethynyl]- (9CI) (CA INDEX NAME)



L6 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:490751 CAPLUS

DOCUMENT NUMBER: 141:63113

TITLE: Macromolecular architectures suitable for  
use in

molecular electronics

INVENTOR(S): Gothelf, Kurt Vesterager; Brown, Raymond S.;  
Thomsen,

Anne; Nielsen, Morten

PATENT ASSIGNEE(S): Aarhus Universitet, Den.

SOURCE: PCT Int. Appl., 139 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.
DATE -----	----	-----	-----
WO 2004050231	A2	20040617	WO 2003-DK821
20031128			
WO 2004050231	A3	20040916	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,			
CA, CH,			
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,			
GB, GD,			
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,			
KZ, LC,			



NI, NO, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
 SY, TJ, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL,  
 ZW TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
 AM, AZ, RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW,  
 DK, EE, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE,  
 SI, SK, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,  
 SN, TD, TG TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,

PRIORITY APPLN. INFO.:  
 20021129

US 2002-429554P P

OTHER SOURCE(S): MARPAT 141:63113

AB A macromol. architecture suitable for use in mol. electronics  
 and in the

manufacture of conductors and semiconductors has been  
 synthesized using linear

and branched oligomers of organic mols. The incorporation of a  
 bi-or

tri-functional organic compound in an oligonucleotide chain and  
 the application

of these for formation of covalently linked organic and  
 metal-organic oligomers

led to a useful mol. architecture. Also, the iterative serial  
 synthesis

of linear and branched organic oligomers by automated methods  
 such as

DNA-synthesis or peptide synthesis, using bi- or tri-functional  
 organic

monomers is described. The compds. may be used to position and  
 arrange

nanoscale substrates such as biomols., biol. structures,  
 colloids,

supramol. structures forming covalently linked assemblies for  
 use as

conducting wires and components in electronic devices. The  
 preparation method

for the macromol. architecture involves the following steps: (1)  
 providing

at least 3 organic compds., each with at least two structural  
 domains with at

least one functional group and an oligonucleotide chain that is  
 at least

partly complementary to the chain on one of the other organic  
 compds.; (2)

hybridizing portions of both oligonucleotide chains in the  
 structural

domains of one compound with one oligonucleotide chains in each  
 of two other


comps.; (3) establishing through the functional groups covalent links between the structural domains joined by the oligonucleotide hybridization; and (4) optionally partly or completely cleaving the oligonucleotide chains formed in step 2.

IT 705930-82-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(macromol. architectures suitable for use in mol. electronics)

RN 705930-82-1 CAPLUS

CN Benzaldehyde, 3,3'-(1,4-phenylenedi-2,1-ethynediyl)bis[5-(1,1-dimethylethyl)-6-hydroxy- (9CI) (CA INDEX NAME)



=>